

Patrick M. Kochanek, MD, MCCM*
 Robert C. Tasker, MA, MD, FRCP*
 Nancy Carney, PhD[§]
 Annette M. Totten, PhD[§]
 P. David Adelson, MD[¶]
 Nathan R. Selden, MD, PhD[¶]
 Cynthia Davis-O'Reilly, BS[§]
 Erica L. Hart, MST[§]
 Michael J. Bell, MD[#]
 Susan L. Bratton, MD, MPH**
 Gerald A. Grant, MD^{##}
 Niranjn Kisson, MD, FRCP(C), MCCM^{§§}
 Karin E. Reuter-Rice, PhD, CPNP-AC^{¶¶}
 Monica S. Vavilala, MD^{|||}
 Mark S. Wainwright, MD, PhD^{##}

*Department of Critical Care Medicine, Department of Anesthesiology, Pediatrics, Bioengineering, and Clinical and Translational Science, Safar Center for Resuscitation Research, University of Pittsburgh School of Medicine, UPMC Children's Hospital of Pittsburgh, Pittsburgh, Pennsylvania; (Continued on next page)

This executive summary has been co-published and appears in *Pediatric Critical Care Medicine* and *Neurosurgery*.

Kochanek PM, Tasker RC, Carney N, Totten AM, Adelson PD, Selden NR, Davis-O'Reilly C, Hart EL, Bell MJ, Bratton SL, Grant GA, Kisson N, Reuter-Rice KE, Vavilala MS, Wainwright MS: Guidelines for the Management of Pediatric Severe Traumatic Brain Injury, Third Edition: Update of the Brain Trauma Foundation Guidelines, Executive Summary. *Pediatr Crit Care Med*. 2019;20(3): 280-289; <https://journals.lww.com/pccmjournals>.

Reviewed for evidence-based integrity and endorsed by the American Association of Neurological Surgeons and Congress of Neurological Surgeons.

Correspondence:

Patrick M. Kochanek, MD, MCCM, Safar Center for Resuscitation Research, UPMC Children's Hospital of Pittsburgh, John G. Rangos Research Center – 6th Floor, 4401 Penn Avenue, Pittsburgh, PA 15224. E-mail: kochanekpm@ccm.upmc.edu

Received, January 11, 2019.

Accepted, February 5, 2019.

Published Online, March 2, 2019.

Copyright © 2019 by the Society of Critical Care Medicine and the World Federation of Pediatric Intensive and Critical Care Societies.

Guidelines for the Management of Pediatric Severe Traumatic Brain Injury, Third Edition: Update of the Brain Trauma Foundation Guidelines, Executive Summary

The purpose of this work is to identify and synthesize research produced since the second edition of these Guidelines was published and incorporate new results into revised evidence-based recommendations for the treatment of severe traumatic brain injury in pediatric patients. This document provides an overview of our process, lists the new research added, and includes the revised recommendations. Recommendations are only provided when there is supporting evidence. This update includes 22 recommendations, 9 are new or revised from previous editions. New recommendations on neuroimaging, hyperosmolar therapy, analgesics and sedatives, seizure prophylaxis, temperature control/hypothermia, and nutrition are provided. None are level I, 3 are level II, and 19 are level III. The Clinical Investigators responsible for these Guidelines also created a companion algorithm that supplements the recommendations with expert consensus where evidence is not available and organizes possible interventions into first and second tier utilization. The complete guideline document and supplemental appendices are available electronically (<https://doi.org/10.1097/PCC.0000000000001735>). The online documents contain summaries and evaluations of all the studies considered, including those from prior editions, and more detailed information on our methodology. New level II and level III evidence-based recommendations and an algorithm provide additional guidance for the development of local protocols to treat pediatric patients with severe traumatic brain injury. Our intention is to identify and institute a sustainable process to update these Guidelines as new evidence becomes available.

KEY WORDS: Critical care, Evidence-based medicine, Head injury, Guidelines, Pediatrics, Systematic review, Traumatic brain injury

Neurosurgery 84:1169–1178, 2019

DOI:10.1093/neuros/nyz051

www.neurosurgery-online.com

The third edition of the Brain Trauma Foundation's Guidelines for the Management of Pediatric Severe Traumatic Brain Injury (TBI)¹ updates the second edition published in 2012.² This new publication is part of an effort to update a suite of 3 Brain Trauma Foundation Guidelines, including similar acute care guidelines for adults (published in January 2017)³ and

guidelines for prehospital management of all ages (forthcoming). It represents a substantial effort by a multidisciplinary group of individuals assembled to reflect the team approach to the treatment of these complex, critically ill patients that is essential to optimizing critical care and improving outcomes.

A total of 48 new studies were included in this third edition. Although some progress has been made and should be celebrated, overall the level of evidence informing these Guidelines remains low. High-quality randomized studies that could support level I recommendations remain absent; the available evidence produced only 3 level II recommendations, whereas most recommendations are level III, supported by lower quality evidence.

ABBREVIATIONS: ADAPT, Aute Pediatric TBI Trial; ACM, advanced cerebral monitoring; CPP, cerebral perfusion pressure; ICP, intracranial pressure; RCT, randomized controlled trial; TBI, Traumatic Brain Injury

Neurosurgery Speaks! Audio abstracts available for this article at www.neurosurgery-online.com.

In addition to the Guidelines, we have authored a companion article that presents a “Critical Pathway” algorithm of care for both first tier and second tier (refractory intracranial hypertension) approaches.⁴ The algorithm reflects both the evidence-based recommendations from these Guidelines as well as consensus-based expert opinion, vetted by the full committee, where evidence was not available. The algorithm also addresses a number of issues that are important but were not previously covered in the Guidelines, given the lack of research. Specifically, the algorithm addresses issues such as a step-wise approach to elevated intracranial pressure (ICP), differences in tempo of therapy in different types of patients, scenarios with a rapidly escalating need for ICP-directed therapy in the setting of impending herniation, integration of multiple monitoring targets, and other complex issues such as minimal vs optimal therapeutic targets and approaches to weaning therapies.

It is important to acknowledge that these Guidelines were written as the Approaches and Decisions in Acute Pediatric TBI Trial (ADAPT),⁵⁻⁷ one of the most important in the field of pediatric TBI, was coming to a close. The ADAPT completed enrollment of 1000 cases of severe pediatric TBI and is one example of the recent, heightened general interest in TBI as a disease. This new interest in the importance of TBI has emerged in part from the recognition of the high prevalence of TBI across the injury severity spectrum, particularly concussion, and from the need for new classification systems and new trial design for TBI in both children and adults.^{8,9} We expect that the results of ADAPT, along with those of other ongoing trials and recently completed research in the field, will help provide new insight and clarity into the acute medical management of infants, children, and adolescents with severe TBI and support further refinement of the recommendations in these documents.

THE SCOPE OF THE GUIDELINES

The Guidelines address monitoring, thresholds for ICP and cerebral perfusion pressure (CPP), and 10 categories of treatments specific to TBI in infants, children, or adolescents. The Guide-

(Continued from previous page)

[†]Department of Neurology, Department of Anesthesiology, Critical Care and Pain Medicine, Boston Children's Hospital; Harvard Medical School, Boston, Massachusetts; [‡]Pacific Northwest Evidence-based Practice Center, Department of Medical Informatics and Clinical Epidemiology, Oregon Health & Science University, Portland, Oregon; [§]Department of Pediatric Neurosurgery, BARROW Neurological Institute at Phoenix Children's Hospital, Phoenix, Arizona; ^{||}Department of Neurological Surgery, Oregon Health & Science University, Portland, Oregon; [¶]Department Critical Care Medicine, Children's National Medical Center, Washington, District of Columbia; ^{**}Department of Pediatrics, University of Utah, Salt Lake City, Utah; ^{††}Department of Neurosurgery, Stanford University, Stanford, California; ^{‡‡}Department of Pediatrics, British Columbia's Children's Hospital, Child and Family Research Institute, University of British Columbia, Vancouver, Canada; ^{§§}School of Nursing/School of Medicine, Department of Pediatrics, Division of Pediatric Critical Care Medicine, Duke University, Durham, North Carolina; ^{||||}Department of Anesthesiology & Pain Medicine, Department of Pediatrics, Harborview Injury Prevention and Research Center (HIPRC), University of Washington, Seattle, Washington; ^{¶¶}Division of Pediatric Neurology, University of Washington, Seattle Children's Hospital, Seattle, Washington

lines are not intended to cover all topics relevant to the care of patients with severe TBI. Specifically, topics related to general good care for all patients, or all trauma patients, are not included.

Developing protocols that integrate TBI-specific, evidence-based recommendations with general best practices for trauma patients, and that provide guidance, suggestions, or options in areas of TBI management where the evidence is insufficient, is outside the scope of these Guidelines. These recommendations are intended to provide the foundation on which protocols can be developed that are appropriate to different treatment environments. The algorithm developed by the clinical investigators is one example of such a protocol, but not the only possible protocol that could be developed based on these Guidelines.

METHODS

The methods for developing these Guidelines were organized in 2 phases—the systematic review, including the identification, assessment, and synthesis of the literature; and the use of that foundation for evidence-based recommendations.

Systematic Evidence Review and Synthesis

Literature Search and Review

Our literature search protocol is described in detail, and the search strategies are in Appendix D of the full online guideline document.¹ Please note that all appendices mentioned in this executive summary refer to appendices to the full guidelines document.¹

The key criteria for including studies in the review were as follows: the population included pediatric patients (age ≤ 18 yr) with severe TBI (defined as Glasgow Coma Scale score of 3-8), and the study assessed an included outcome (mortality, neurologic function, or appropriate intermediate outcomes for the topic). Two reviewers independently identified studies to include, and differences were resolved via consensus or by a third reviewer. Detailed inclusion criteria and a list of studies excluded after full-text review are in the online document in Appendices B and E.¹ This edition adds studies published from 2010 to June 2017.

Quality Assessment and Data Abstraction of Individual Studies

All included studies were assessed for potential for bias, which is a systematic approach to assessing the internal validity or quality of studies. The quality criteria used in the second edition were maintained and applied to the newly identified studies of monitoring and treatments. The criteria for threshold studies were revised to be specific to the quality of threshold studies. (See appendix F in the online document¹ for a complete list of the quality criteria used for individual studies.) Key data elements were then extracted from each study and placed into tables. The tables were provided to the clinical investigators and summarized by topic in the guideline document (see summaries by topic in the full report online).¹ Class 1 is the highest class and is limited to good-quality randomized trials. Class 2 includes moderate-quality randomized controlled trials (RCTs) and good-quality cohort or case-control studies. Class 3 is the lowest class and is given to low-quality RCTs, moderate to low-quality cohort or case-control studies, and treatment series and other noncomparative designs.

Synthesis

The final phase of the evidence review is the synthesis of individual studies into information that the clinical investigators and the methods team use to develop recommendations. This synthesis is described for each topic in the online document in the sections titled evaluation of the evidence, following the recommendations and preceding the evidence summary.

Quality of the Body of Evidence

Assessing the quality of the body of evidence involves four domains: the aggregate quality of the studies, the consistency of the results, whether the evidence provided is direct or indirect, and the precision of the effect estimates. The criteria and ratings are outlined in the Methods section of the online document and more detailed definitions are in Appendix G.¹ In addition, the number of studies and number of included subjects are considered. Based on these, an overall assessment is made as to whether the quality of the body of evidence is high, moderate, low, or insufficient. The assessment of the body of evidence for each subtopic is included in a table in each topic section in the full guideline document.

Applicability

Applicability is the extent to which research findings are useful for informing recommendations for a broader population (usually the population that is the target of the recommendations). In this edition, we considered the applicability of individual studies in the quality of the body of evidence and applicability section immediately following the recommendations in the full guideline document.

Recommendations

Development of Recommendations

Classes 1, 2, and 3 studies constitute the evidence on which the recommendations are based. Once evidence was identified, whether or not it could be used to inform recommendations was based on the quality of the body of evidence and consideration of applicability. Under our current methods, identification of evidence is necessary but not sufficient for the development of evidence-based recommendations. If no evidence was identified, no recommendations were made. If the identified evidence was extremely limited (eg, inconsistent results, imprecise), it could be considered insufficient to support a recommendation.

Given this approach, there were cases in which evidence was identified, but the quality was low and applicability concerns restricted the ability to translate the evidence into recommendations. Even if a recommendation was not made, the studies contributing evidence were included in the full Guideline to acknowledge their place in the body of evidence and make the evidence accessible for future consideration. As new studies are generated and added to the evidence base, we expect to see changes in the assessment of the quality of the body of evidence.

Level of Recommendations

Recommendations in this edition are designated as level I, level II, or level III. The level of recommendation is determined by the assessment of the quality of the body of evidence, which includes, but is not limited to, the class of the included studies.

The levels were primarily based on the quality of the body of evidence as follows:

- 1) Level I recommendations were based on a high-quality body of evidence.

- 2) Level II recommendations were based on a moderate-quality body of evidence.
- 3) Level III recommendations were based on a low-quality body of evidence.

In addition to the quality of evidence, we also considered applicability. Currently, there is a lack of standards and developed methods to assess applicability. For this reason, applicability alone was not used to downgrade a recommendation; however, we did include and document in the full guideline any applicability issues that were identified and discussed by the authors.

“Insufficient” was used in cases where the body of evidence was insufficient to support a recommendation either because there were no studies identified or because the body of evidence had major quality limitations. If the evidence was rated insufficient, no recommendation was made.

REVISED RECOMMENDATIONS

Summary of Changes to Recommendations

This update includes 22 evidence-based recommendations; 9 are new or revised significantly from the previous edition. There are no level I recommendations, 3 recommendations are level II, and the remaining 19 are level III.

Tables 1, 2, and 3 provide the recommendations for monitoring, thresholds, and treatments, respectively. Each recommendation is numbered with a roman numeral for the level followed by a period and a number counting the recommendations in each topic (So III.1 is the first Level III recommendation and III.2 is the second level III recommendation). In these tables, the recommendations in bold are new or have been significantly revised, whereas those in regular text have not changed or only have changes in wording. The online guideline document includes a section on each topic consisting of an Introduction, Recommendations, Evaluation of the Evidence, and Summary of the Evidence (including evidence tables and a narrative overview).

Monitoring Recommendations

Monitoring does not affect outcomes directly; rather the information from monitoring can be used to direct treatment decisions. Treatment informed by data from monitoring may result in better outcomes than treatment informed solely by data from clinical assessment. Monitoring recommendations are related to the influence on patient outcomes of three types of monitoring: ICP monitoring, advanced cerebral monitoring (ACM), and neuroimaging. The recommendations for ICP and ACM did not change; however, two notes were added to the ACM recommendation. For neuroimaging, one new recommendation suggesting that CT examinations not be used to rule out the possibility of elevated ICP was added to the existing recommendation.

Threshold Recommendations

These recommendations are related to threshold values for variables that are monitored during the in-hospital management of patients with severe TBI. This includes thresholds for ICP and CPP. There are no changes to the recommendations from

TABLE 1. Updated Recommendations: Monitoring	
Topics	Recommendations
Intracranial pressure monitoring	Level III To improve overall outcomes III.1. Use of ICP monitoring is suggested.
Advanced neuromonitoring	Level III To improve overall outcomes III.1. If PbrO ₂ monitoring is used, maintaining a level >10 mm Hg is suggested. <i>Note 1.</i> There was insufficient evidence to support a recommendation for the use of a monitor of Po ₂ in brain interstitium (PbrO ₂) to improve outcomes. <i>Note 2.</i> Use of advanced neuromonitoring (brain oxygenation) should only be for patients with no contraindications to invasive neuromonitoring, such as coagulopathy, and for patients who do not have a diagnosis of brain death.
Neuroimaging	Level III To improve overall outcomes III.1. Excluding the possibility of elevated ICP on the basis of a normal initial (0-6 h after injury) CT examination of the brain is not suggested in comatose pediatric patients. III.2. Routinely obtaining a repeat CT scan >24 hours after the admission and initial follow-up is not suggested for decisions about neurosurgical intervention, unless there is either evidence of neurologic deterioration or increasing ICP.

ICP = intracranial pressure, PbrO₂ = brain tissue oxygen.
Bold indicates new or revised recommendations.

TABLE 2. Updated Recommendations: Thresholds	
Topics	Recommendations
Threshold for treatment of intracranial hypertension	Level III To improve overall outcomes III.1. Treatment of intracranial pressure targeting a threshold of < 20 mm Hg is suggested.
Thresholds for cerebral perfusion pressure	Level III To improve overall outcomes III.1 Treatment to maintain CPP at a minimum of 40 mm Hg is suggested. III.2. A CPP target between 40 and 50 mm Hg is suggested to ensure that the minimum value of 40 mm Hg is not breached. There may be age-specific thresholds with infants at the lower end and adolescents at or above the upper end of this range.

CPP = cerebral perfusion pressure.

the prior edition. Additional studies that supported the existing recommendations were added to the evidence tables in the full guideline document and are listed in Table 4.

Treatment Recommendations

Table 3 contains the recommendations for 10 treatments included in the Guidelines. These topics are included because they are specific to the in-hospital management of TBI or are related to risks experienced by pediatric TBI patients. The topics that are included reflect current practice but are expected to change as new treatments are developed that may replace or complement existing treatments. These topics include 15 recommendations; of these 7 are new or revised. These 7 include 2 recommendations in hyperosmolar therapy; 1 in analgesics, sedatives, and neuromuscular blockade; 1 in seizure prophylaxis; 2 in temperature control; and 1 in nutrition.

DISCUSSION

New Evidence

Table 4 lists the 35 new studies¹⁰⁻⁴⁶ added to the evidence base that was used to support new or existing recommendations. This table presents the studies by topic, provides the citation, and includes the study design, the number of patients included (n), and data class. An additional 13 new studies were added to the guideline document that addressed topics without sufficient evidence to support a recommendation.⁴⁷⁻⁵⁹ More details, such as the outcomes and results for all new studies, are included in the evidence tables and narrative in the full online guideline.

Ongoing and Future Research

Evidence-based guidelines rarely (if ever) contain enough data to fully populate a clinical protocol. This is certainly the case

TABLE 3. Updated Recommendations: Treatments

Topics	Recommendations
Hyperosmolar therapy	<p>Level II For ICP control II.1. Bolus hypertonic saline (3%) is recommended in patients with intracranial hypertension. Recommended effective doses for acute use range between 2 and 5 mL/kg over 10-20 min.</p> <p>Level III For ICP control III.1. Continuous infusion hypertonic saline is suggested in patients with intracranial hypertension. Suggested effective doses as a continuous infusion of 3% saline range from between 0.1 and 1.0 mL/kg of body weight per hour, administered on a sliding scale. The minimum dose needed to maintain intracranial pressure ICP < 20 mm Hg is suggested. III.2. Bolus of 23.4% hypertonic saline is suggested for refractory ICP. The suggested dose is 0.5 mL/kg with a maximum of 30 mL.</p> <p><u>Safety recommendation</u> (applies to all recommendations for this topic). In the context of multiple ICP related therapies, avoiding sustained (>72 h) serum sodium >170 mEq/L is suggested to avoid complications of thrombocytopenia and anemia, whereas avoiding a sustained serum sodium >160 mEq/L is suggested to avoid the complication of deep vein thrombosis.</p> <p><u>Note.</u> Although mannitol is commonly used in the management of raised ICP in pediatric traumatic brain injury, no studies meeting inclusion criteria were identified for use as evidence for this topic.</p>
Analgesics, sedatives, and neuromuscular blockade	<p>Level III For ICP control III.1. With use of multiple ICP-related therapies, as well as appropriate use of analgesia and sedation in routine ICU care, avoiding bolus administration of midazolam and/or fentanyl during ICP crises is suggested due to risks of cerebral hypoperfusion.</p> <p><u>Note 1.</u> In the absence of outcome data, the specific indications, choice, and dosing of analgesics, sedatives, and neuromuscular blocking agents should be left to the treating physician.</p> <p><u>Note 2.</u> Based on guidance from the US Food and Drug Administration, prolonged continuous infusion of propofol for either sedation or the management of refractory intracranial hypertension is not recommended.</p>
Cerebrospinal fluid drainage	<p>Level III For ICP control III.1. Cerebrospinal fluid drainage through an external ventricular drain is suggested to manage increased ICP.</p>
Seizure prophylaxis	<p>Level III For seizure prevention (clinical and subclinical) III.1. Prophylactic treatment is suggested to reduce the incidence of early (within 7 d) PTS.</p> <p><u>Note.</u> At the present time there is insufficient evidence to recommend levetiracetam over phenytoin based on either efficacy in preventing early PTS or toxicity.</p>
Ventilation therapies	<p>Level III To improve overall outcomes III.1. Prophylactic severe hyperventilation to a PaCO₂ < 30 mm Hg in the initial 48 h after injury is not suggested. III.2. If hyperventilation is used in the management of refractory intracranial hypertension, advanced neuromonitoring for evaluation of cerebral ischemia is suggested.</p>
Temperature control/hypothermia*	<p>Level II To improve overall outcomes II.1. Prophylactic moderate (32° C to 33° C) hypothermia is not recommended over normothermia to improve overall outcomes.</p> <p>Level III For ICP control III.1. Moderate (32° C to 33° C) hypothermia is suggested for ICP control.</p> <p><u>Safety recommendation 1.</u> If hypothermia is used and rewarming is initiated, it should be carried out at a rate of 0.5° C to 1.0° C every 12-24 h or slower to avoid complications.</p> <p><u>Safety recommendation 2.</u> If phenytoin is used during hypothermia, monitoring and dosing adjusted to minimize toxicity, especially during the rewarming period, is suggested.</p>

TABLE 3. Continued

Topics	Recommendations
Barbiturates	<p>Level III For ICP control III.1. High-dose barbiturate therapy is suggested in hemodynamically stable patients with refractory intracranial hypertension despite maximal medical and surgical management. <u>Safety recommendation.</u> When high-dose barbiturate therapy is used to treat refractory intracranial hypertension, continuous arterial blood pressure monitoring and cardiovascular support to maintain adequate cerebral perfusion pressure are required because cardiorespiratory instability is common among patients treated with barbiturate coma.</p>
Decompressive craniectomy	<p>Level III For ICP control III.1. Decompressive craniectomy is suggested to treat neurologic deterioration, herniation, or intracranial hypertension refractory to medical management.</p>
Nutrition	<p>Level II To improve overall outcomes II.1. Use of an immune-modulating diet is not recommended.</p> <p>Level III To improve overall outcomes III.1. Initiation of early enteral nutritional support (within 72 h from injury) is suggested to decrease mortality and improve outcomes.</p>
Corticosteroids	<p>Level III To improve overall outcomes III.1. The use of corticosteroids is not suggested to improve outcome or reduce ICP. <u>Note.</u> Recommendation III.1 is not intended to circumvent use of replacement corticosteroids for patients needing chronic steroid replacement therapy, those with adrenal suppression, and those with injury to the hypothalamic-pituitary steroid axis.</p>

Bold: New or revised recommendations.

ICP = intracranial Pressure, PTS = posttraumatic seizures.

*The first recommendation indicates that prophylactic hypothermia does not improve overall outcomes for pediatric patients with severe TBI. The second recommendation indicates that hypothermia is effective in control of ICP. Although this may appear to be somewhat antithetical, the 2 endpoints of overall outcomes and ICP control are clearly distinct. Please see the full Guideline for additional detail.

with the treatment of severe pediatric TBI. Rather the goal is to contribute to a transparent, ongoing process that leads to better research and more evidence in the future. These Guidelines provide recommendations based on the available evidence and at the same time identify gaps that can inform the future research agenda. These gaps can be filled by creating clinical protocols using consensus where evidence is lacking. Together the gaps and protocols provide structure and identify patient samples for the generation of new research. The new research populates the evidence base which can then be used to further develop the Guidelines, creating a recursive cycle designed to grow the evidence base and increase the number of evidence-based recommendations in the future.

Although the number of studies has increased in this update of the Guidelines, most recommendations are based on a small number of studies that are mostly class 3. We hope this will change as the impact of evidence-based practice is documented and new studies undertaken. We are optimistic that the next update will have a stronger evidence base because an important study of pediatric TBI, designed and executed by a guidelines clinical investigator and coauthor, is concluding. This study, ADAPT, was designed to address 12 a priori hypotheses across five Guidelines topics (advanced neuromonitoring, hyperosmolar therapy, cerebrospinal fluid drainage, ventilation, and nutrition) and is

likely to also provide information on other topics and questions from post hoc analyses.⁵ ADAPT is an important example of the value of a guideline in highlighting what cannot be said due to lack of evidence; those gaps provide opportunities for innovation and direction for research.

In addition to ADAPT, the pediatric TBI community needs to promote and support innovative ways to generate higher quality class 1 and class 2 studies that can inform stronger (ie, level I and level II) recommendations. These other needs include the following:

- 1) Research that examines the integration of individual treatments in the context of goal-directed therapy. No treatment or management approach exists independent of other treatments and approaches or independent of the ecology of the treatment setting. The design of meaningful and effective future research must be consistent with this clinical reality.
- 2) Ongoing identification of new topics for investigation. As our understanding of TBI and trauma improves, it is likely new topics will need to be added to the Guidelines. The literature and ongoing trials need to be scanned regularly. It is important that the Guidelines reliably include what evidence is available for new, emerging topics and treatments.

TABLE 4. New Studies Added Since Last Edition to Evidence Supporting Revisions to Recommendations

Topic	Reference	Study design and sample size (n)	Data class	
Monitoring				
Intracranial pressure monitoring	Bennett et al ¹⁰	Retrospective, n = 3084	3	
	Alkoury et al ¹¹	Retrospective, n = 3107	3	
	Bennett et al ¹²	Retrospective, n = 4667	3	
Advanced neuromonitoring	Stippler et al ¹³	Treatment series, n = 46	3	
	Figaji et al ¹⁴	Treatment series, n = 28	3	
Neuroimaging	Bailey et al ¹⁵	Treatment series, n = 9	3	
	Bata, 2014 et al ¹⁶	Retrospective, n = 71	3	
Thresholds				
Thresholds for treatment of intracranial hypertension	Miller Ferguson et al ¹⁷	Retrospective, n = 85	3	
	Mehta et al ¹⁸	Retrospective, n = 22	3	
Thresholds for cerebral perfusion pressure	Allen et al ¹⁹	Retrospective, n = 317	2	
	Miller Ferguson et al ¹⁷	Retrospective, n = 85	3	
	Vavilala et al ²⁰	Retrospective, n = 236	3	
Treatments				
Hyperosmolar therapy	Shein et al ²¹	Prospective, n = 16	2	
	Piper et al ²²	Treatment series, n = 32	3	
	Webster et al ²³	Retrospective, n = 58	3	
	Gonda et al ²⁴	Retrospective, n = 48 traumatic brain injury	3	
Analgesics, sedatives, and neuromuscular blockade	Welch et al ²⁵	Treatment series, n = 31	3	
Cerebrospinal fluid drainage	Shein et al ²¹	Prospective, n = 16	3	
	Andrade et al ²⁶	Treatment series, n = 58, n = 11 (younger than 17)	3	
Seizure prophylaxis	Liesemer et al ²⁷	Retrospective, n = 54 moderate, n = 221 severe	3	
Ventilation therapies	No New Studies			
Temperature control/hypothermia	Tasker et al ²⁸	Meta-analysis, n = 470	Fair quality	
	Crompton et al ²⁹⁻³¹	Meta-analysis, n = 454	Poor quality	
	Adelson et al ³²	RCT, n = 77	1	
	Beca et al ³³	RCT, n = 50	2	
	Hutchinson et al ³⁴	Retrospective (secondary analysis of 2008 RCT), n = 225	2	
	Empey et al ³⁵	RCT, n = 19	3	
	Vavilala et al ²⁰	Retrospective, n = 236	3	
Barbiturates	Mellion et al ³⁶	Treatment series, n = 36	3	
	Pechmann et al ³⁷	Treatment series, n = 12	3	
Decompressive craniectomy	Prasad et al ³⁸	Treatment series, n = 71	3	
	Desgranges et al ³⁹	Treatment series, n = 12	3	
	Khan et al ⁴⁰	Treatment series, n = 25, 21 severe	3	
	Csokay et al ⁴¹	Treatment series, n = 8	3	
	Suarez et al ⁴²	Treatment series, n = 14	3	
	Adamo et al ⁴³	Treatment series, n = 7	3	
	Figaji et al ⁴⁴	Treatment series, n = 12	3	
	Messing-Junger et al ⁴⁵	Treatment series, n = 7	3	
	Nutrition	Taha et al ⁴⁶	Retrospective, n = 90	3
	Corticosteroids	No New Studies		

RCT = randomized controlled trial.

3) Consistency in data collection across studies. Future research should emphasize consistency in data collection across research projects, such as utilization of the common data elements of the National Institutes of Health.⁶⁰⁻⁶³

It is important that the pediatric TBI research community systematically address these questions by creating a prioritized research agenda and advocating for additional high-quality research that can populate the evidence base for future guidelines.

CONCLUSION

The increase in the number of studies as well as the number of class 2 studies and level II recommendations is encouraging. The growth in the evidence base strengthens the utility of the evidence-based recommendations as a basis for local protocols, which can incorporate consensus where evidence is still not available. However, this update also underscores that much work remains to be done if our goal is evidence-based treatment designed to improve outcomes for children who sustain severe TBI.

Disclaimer

Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the U.S. Army Contracting Command, Aberdeen Proving Ground, Natick Contracting Division, Stanford University, or the Brain Trauma Foundation. The information contained in the Guidelines for the Management of Pediatric Severe TBI reflects the current state of knowledge at the time of publication. The Brain Trauma Foundation, American Association of Neurological Surgeons, Congress of Neurological Surgeons, and other collaborating organizations are not engaged in rendering professional medical services and assume no responsibility for patient outcomes resulting from application of these general recommendations in specific patient circumstances. Accordingly, the Brain Trauma Foundation, American Association of Neurological Surgeons, and Congress of Neurological Surgeons consider adherence to these clinical practice guidelines will not necessarily assure a successful medical outcome. The information contained in these guidelines reflects published scientific evidence at the time of completion of the guidelines and cannot anticipate subsequent findings and/or additional evidence and therefore should not be considered inclusive of all proper procedures and tests or exclusive of other procedures and tests that are reasonably directed to obtaining the same result. Medical advice and decisions are appropriately made only by a competent and licensed physician who must make decisions in light of all the facts and circumstances in each individual and particular case and on the basis of availability of resources and expertise. Guidelines are not intended to supplant physician judgment with respect to particular patients or special clinical situations and are not a substitute for physician-patient consultation. Accordingly, the Brain Trauma Foundation, American Association of Neurological Surgeons, and Congress of Neurological Surgeons consider adherence to these guidelines to be voluntary, with the ultimate determination regarding their application to be made by the physician in light of each patient's individual circumstances.

Funding

Supported, in part, by the U.S. Army Contracting Command, Aberdeen Proving Ground, Natick Contracting Division, through a contract awarded to Stanford University (W911 QY-14-C-0086) and a subcontract awarded to Oregon Health & Science University. Prior editions were supported, in part, by funding from multiple sources through the Brain Trauma Foundation. Dr Kochanek received funding from the Society of Critical Care Medicine (Editor-in-Chief of Pediatric Critical Care Medicine), from serving as an expert witness on cases in

pediatric critical care. Drs Carney and Totten's, Ms Davis-O'Reilly's, and Ms Hart's institutions received funding from Stanford University. Dr Selden disclosed that he has stock options (current \$0 value) in Cerebrotech for scientific advisory board service (this device is not clinically available and is not referenced in the work). Dr Reuter-Rice received funding from textbook royalties and curriculum content, and she received support for article research from Robert Wood Johnson Foundation funding 2013–2016. Dr Wainwright received funding from Sage Therapeutics. The remaining authors have disclosed that they do not have any potential conflicts of interest.

REFERENCES

- Kochanek PM, Tasker RC, Carney N, et al. Guidelines for the management of pediatric severe traumatic brain injury, third edition: update of the brain trauma foundation guidelines. *Pediatr Crit Care Med.* 2019;20(Suppl 1): S1-S82.
- Kochanek PM, Carney N, Adelson PD, et al. Guidelines for the acute medical management of severe traumatic brain injury in infants, children, and adolescents—second edition. *Pediatr Crit Care Med.* 2012;13(Suppl 1): S1-S82.
- Carney N, Totten AM, O'Reilly C, et al. Guidelines for the management of severe traumatic brain injury, fourth edition. *Neurosurgery.* 2017;80(1):6-15.
- Kochanek PM, Tasker RC, Bell MJ, et al. Pediatric Severe Traumatic Brain Injury: 2019 Consensus and Guidelines-Based Algorithm for First and Second Tier Therapies. *Pediatr Crit Care Med.* 2019;20(3):269-279.
- Bell MJ, Adelson PD, Hutchison JS, et al. Differences in medical therapy goals for children with severe traumatic brain injury—an international study. *Pediatr Crit Care Med.* 2013;14(8):811-818.
- Bell MJ, Adelson PD, Wisniewski SR. Challenges and opportunities for pediatric severe TBI—review of the evidence and exploring a way forward. *Childs Nerv Syst.* 2017;33(10):1663-1667.
- Kurz JE, Poloyac SM, Abend NS, Fabio A, Bell MJ, Wainwright MS. Variation in anticonvulsant selection and electroencephalographic monitoring following severe traumatic brain injury in children—understanding resource availability in sites participating in a comparative effectiveness study. *Pediatr Crit Care Med.* 2016;17(7):649-657.
- Jha RM, Kochanek PM. Adding insight to injury: a new era in neurotrauma. *Lancet Neurol.* 2017;16(8):578-580.
- Kochanek PM, Bell MJ. Tackling the challenges of clinical trials for severe traumatic brain injury in children: screening, phenotyping, and adapting. *Crit Care Med.* 2015;43(7):1544-1546.
- Bennett TD, DeWitt PE, Greene TH, et al. Functional outcome after intracranial pressure monitoring for children with severe traumatic brain injury. *JAMA Pediatr.* 2017;171(10):965-971.
- Alkhoury F, Kyriakides TC. Intracranial pressure monitoring in children with severe traumatic brain injury: National Trauma Data Bank-based review of outcomes. *JAMA Surg.* 2014;149(6):544-548.
- Bennett TD, Riva-Cambrin J, Keenan HT, Korgenski EK, Bratton SL. Variation in intracranial pressure monitoring and outcomes in pediatric traumatic brain injury. *Arch Pediatr Adolesc Med.* 2012;166(7):641-647.
- Stippler M, Ortiz V, Adelson PD, et al. Brain tissue oxygen monitoring after severe traumatic brain injury in children: relationship to outcome and association with other clinical parameters. *J Neurosurg Pediatr.* 2012;10(5):383-391.
- Figaji AA, Zwane E, Graham Fieggen A, Argent AC, Le Roux PD, Peter JC. The effect of increased inspired fraction of oxygen on brain tissue oxygen tension in children with severe traumatic brain injury. *Neurocrit Care.* 2010;12(3): 430-437.
- Bailey BM, Liesemer K, Statler KD, Riva-Cambrin J, Bratton SL. Monitoring and prediction of intracranial hypertension in pediatric traumatic brain injury: clinical factors and initial head computed tomography. *J Trauma Acute Care Surg.* 2012;72(1):263-270.
- Bata SC, Yung M. Role of routine repeat head imaging in paediatric traumatic brain injury. *ANZ J Surg.* 2014;84(6):438-441.
- Miller Ferguson N, Shein SL, Kochanek PM, et al. Intracranial hypertension and cerebral hypoperfusion in children with severe traumatic brain injury: thresholds and burden in accidental and abusive insults. *Pediatr Crit Care Med.* 2016;17(5):444-450.

18. Mehta A, Kochanek PM, Tyler-Kabara E, et al. Relationship of intracranial pressure and cerebral perfusion pressure with outcome in young children after severe traumatic brain injury. *Dev Neurosci*. 2010;32(5-6):413-419.
19. Allen BB, Chiu YL, Gerber LM, Ghajar J, Greenfield JP. Age-specific cerebral perfusion pressure thresholds and survival in children and adolescents with severe traumatic brain injury. *Pediatr Crit Care Med*. 2014;15(1):62-70.
20. Vavilala MS, Kernic MA, Wang J, et al. Acute care clinical indicators associated with discharge outcomes in children with severe traumatic brain injury. *Crit Care Med*. 2014;42(10):2258-2266.
21. Shein SL, Ferguson NM, Kochanek PM, et al. Effectiveness of pharmacological therapies for intracranial hypertension in children with severe traumatic brain injury—results from an automated data collection system time-synched to drug administration. *Pediatr Crit Care Med*. 2016;17(3):236-245.
22. Piper BJ, Harrigan PW. Hypertonic saline in paediatric traumatic brain injury: a review of nine years' experience with 23.4% hypertonic saline as standard hyperosmolar therapy. *Anaesth Intensive Care*. 2015;43(2):204-210.
23. Webster DL, Fei L, Falcone RA, Kaplan JM. Higher-volume hypertonic saline and increased thrombotic risk in pediatric traumatic brain injury. *J Crit Care*. 2015;30(6):1267-1271.
24. Gonda DD, Meltzer HS, Crawford JR, et al. Complications associated with prolonged hypertonic saline therapy in children with elevated intracranial pressure. *Pediatr Crit Care Med*. 2013;14(6):610-620.
25. Welch TP, Wallendorf MJ, Kharasch ED, Leonard JR, Doctor A, Pineda JA. Fentanyl and midazolam are ineffective in reducing episodic intracranial hypertension in severe pediatric traumatic brain injury. *Crit Care Med*. 2016;44(4):809-818.
26. Andrade AF, Paiva WS, Amorim RL, et al. Continuous ventricular cerebrospinal fluid drainage with intracranial pressure monitoring for management of posttraumatic diffuse brain swelling. *Arq Neuro-Psiquiatr*. 2011;69(1):79-84.
27. Liesemer K, Bratton SL, Zebrock CM, Brockmeyer D, Statler KD. Early post-traumatic seizures in moderate to severe pediatric traumatic brain injury: rates, risk factors, and clinical features. *J Neurotrauma*. 2011;28(5):755-762.
28. Tasker RC, Vonberg FW, Ulano ED, Akhondi-Asl A. Updating evidence for using hypothermia in pediatric severe traumatic brain injury: conventional and Bayesian meta-analytic perspectives. *Pediatr Crit Care Med*. 2017;18(4):355-362.
29. Crompton EM, Lubomirova I, Cotlarciuc I, Han TS, Sharma SD, Sharma P. Meta-analysis of the therapeutic hypothermia for traumatic brain injury in adult and pediatric patients. *Crit Care Med*. 2017;45(4):575-583.
30. Tasker RC, Akhondi-Asl A. Updating evidence for using therapeutic hypothermia in pediatric severe traumatic brain injury. *Crit Care Med*. 2017;45(10):e1091.
31. Crompton E, Sharma P. The authors reply. *Crit Care Med*. 2017;45(10):e1091-e1092.
32. Adelson PD, Wisniewski SR, Beca J, et al. Comparison of hypothermia and normothermia after severe traumatic brain injury in children (Cool Kids): a phase 3, randomised controlled trial. *Lancet Neurol*. 2013;12(6):546-553.
33. Beca J, McSharry B, Erickson S, et al. Hypothermia for traumatic brain injury in children—a phase II randomized controlled trial. *Crit Care Med*. 2015;43(7):1458-1466.
34. Hutchison JS, Frndova H, Lo TY, Guerguerian AM, Hypothermia Pediatric Head Injury Trial Investigators Canadian Critical Care Trials Group. Impact of hypotension and low cerebral perfusion pressure on outcomes in children treated with hypothermia therapy following severe traumatic brain injury: a post hoc analysis of the Hypothermia Pediatric Head Injury Trial. *Dev Neurosci*. 2011;32(5-6):406-412.
35. Empey PE, de Mendizabal NV, Bell MJ, et al. Therapeutic hypothermia decreases phenytoin elimination in children with traumatic brain injury. *Crit Care Med*. 2013;41(10):2379-2387.
36. Mellion SA, Bennett KS, Ellsworth GL, et al. High-dose barbiturates for refractory intracranial hypertension in children with severe traumatic brain injury. *Pediatr Crit Care Med*. 2013;14(3):239-247.
37. Pechmann A, Anastasopoulos K, Korinthenberg R, van Velthoven-Wurster V, Kirschner J. Decompressive craniectomy after severe traumatic brain injury in children: complications and outcome. *Neuropediatrics*. 2015;46(01):005-012.
38. Prasad GL, Gupta DK, Mahapatra AK, Sharma BS. Surgical results of decompressive craniectomy in very young children: a level one trauma centre experience from India. *Brain Inj*. 2015;29(13-14):1717-1724.
39. Desgranges FP, Javouhey E, Mottolese C, et al. Intraoperative blood loss during decompressive craniectomy for intractable intracranial hypertension after severe traumatic brain injury in children. *Childs Nerv Syst*. 2014;30(8):1393-1398.
40. Khan SA, Shallwani H, Shamim MS, et al. Predictors of poor outcome of decompressive craniectomy in pediatric patients with severe traumatic brain injury: a retrospective single center study from Pakistan. *Childs Nerv Syst*. 2014;30(2):277-281.
41. Csokay A, Emelifeonwu JA, Fugedi L, Valalik I, Lang J. The importance of very early decompressive craniectomy as a prevention to avoid the sudden increase of intracranial pressure in children with severe traumatic brain swelling (retrospective case series). *Childs Nerv Syst*. 2012;28(3):441-444.
42. Suarez EP, Gonzalez AS, Diaz CP, Salido AG, de Azagra Garde AM, Flores JC. Decompressive craniectomy in 14 children with severe head injury: clinical results with long-term follow-up and review of the literature. *J Trauma*. 2011;71(1):133-140.
43. Adamo MA, Drazin D, Waldman JB. Decompressive craniectomy and postoperative complication management in infants and toddlers with severe traumatic brain injuries. *J Neurosurg Pediatr*. 2009;3(4):334-339.
44. Figaji AA, Fieggen AG, Argent A, Peter JC. Surgical treatment for “brain compartment syndrome” in children with severe head injury. *S Afr Med J*. 2006;96(9 Pt 2):969-975.
45. Messing-Junger AM, Marzog J, Wobker G, Sabel M, Bock WJ. Decompressive craniectomy in severe brain injury. *Zentralbl Neurochir*. 2003;64(04):171-177.
46. Taha AA, Badr L, Westlake C, Dec V, Mudit M, Tiras KL. Effect of early nutritional support on intensive care unit length of stay and neurological status at discharge in children with severe traumatic brain injury. *J Neurosci Nurs*. 2011;43(6):291-297.
47. Bar-Joseph G, Guilburd Y, Tamir A, Guilburd JN. Effectiveness of ketamine in decreasing intracranial pressure in children with intracranial hypertension. *J Neurosurg Pediatr*. 2009;4(1):40-46.
48. Bourdages M, Bigras J-L, Farrell CA, Hutchison JS, Lacroix J, Canadian Critical Care Trials G. Cardiac arrhythmias associated with severe traumatic brain injury and hypothermia therapy. *Pediatr Crit Care Med*. 2010;11(3):408-414.
49. Chin KH, Bell MJ, Wisniewski SR, et al. Effect of administration of neuromuscular blocking agents in children with severe traumatic brain injury on acute complication rates and outcomes: a secondary analysis from a randomized, controlled trial of therapeutic hypothermia. *Pediatr Crit Care Med*. 2015;16(4):352-358.
50. Chung MG, O'Brien NF. Prevalence of early posttraumatic seizures in children with moderate to severe traumatic brain injury despite levetiracetam prophylaxis. *Pediatr Crit Care Med*. 2016;17(2):150-156.
51. Josan VA, Sgouros S. Early decompressive craniectomy may be effective in the treatment of refractory intracranial hypertension after traumatic brain injury. *Childs Nerv Syst*. 2006;22(10):1268-1274.
52. Mhanna MJ, Mallah WE, Verrees M, Shah R, Super DM. Outcome of children with severe traumatic brain injury who are treated with decompressive craniectomy. *J Neurosurg Pediatr*. 2015;16(5):508-514.
53. Oluigbo CO, Wilkinson CC, Stence NV, Fenton LZ, McNatt SA, Handler MH. Comparison of outcomes following decompressive craniectomy in children with accidental and nonaccidental blunt cranial trauma. *J Neurosurg Pediatr*. 2012;9(2):125-132.
54. Pearl PL, McCarter R, McGavin CL, et al. Results of phase II levetiracetam trial following acute head injury in children at risk for posttraumatic epilepsy. *Epilepsia*. 2013;54(9):e135-e137.
55. Roumeliotis N, Dong C, Pettersen G, Crevier L, Emeriaud G. Hyperosmolar therapy in pediatric traumatic brain injury: a retrospective study. *Childs Nerv Syst*. 2016;32(12):2363-2368.
56. Rubiano AM, Villarreal W, Hakim EJ, et al. Early decompressive craniectomy for neurotrauma: an institutional experience. *Ulus Travma Acil Cerrahi Derg*. 2009;15(1):28-38.
57. Su E, Bell MJ, Kochanek PM, et al. Increased CSF concentrations of myelin basic protein after TBI in infants and children: absence of significant effect of therapeutic hypothermia. *Neurocrit Care*. 2012;17(3):401-407.
58. Taylor A, Butt W, Rosenfeld J, et al. A randomized trial of very early decompressive craniectomy in children with traumatic brain injury and sustained intracranial hypertension. *Child's Nervous System*. 2001;17(3):154-162.

59. Thomale U-W, Graetz D, Vajkoczy P, Sarrafzadeh AS. Severe traumatic brain injury in children—a single center experience regarding therapy and long-term outcome. *Childs Nerv Syst.* 2010;26(11):1563-1573.
60. Adelson PD, Pineda J, Bell MJ, et al. Common data elements for pediatric traumatic brain injury: recommendations from the working group on demographics and clinical assessment. *J Neurotrauma.* 2012;29(4):639-653.
61. Berger RP, Beers SR, Papa L, Bell M. Common data elements for pediatric traumatic brain injury: recommendations from the biospecimens and biomarkers workgroup. *J Neurotrauma.* 2012;29(4):672-677.
62. Duhaime AC, Holsouser B, Hunter JV, Tong K. Common data elements for neuroimaging of traumatic brain injury: pediatric considerations. *J Neurotrauma.* 2012;29(4):629-633.
63. McCauley SR, Wilde EA, Anderson VA, et al. Recommendations for the use of common outcome measures in pediatric traumatic brain injury research. *J Neurotrauma.* 2012;29(4):678-705.

Acknowledgments

We would like to thank the following people at the Pacific Northwest Evidence-based Practice Center at Oregon Health & Science University for their invaluable assistance in producing this document: Roger Chou, MD, Elaine Graham, MLS, Andrew Hamilton, MS, MLS, Hyon Hildebrant, BA, Shaun Ramirez, MPH, Leah Williams, BS. We also thank Jamshid Ghajar, MD, PhD, from the Brain Trauma Foundation and Stanford University.

Neurosurgery Speaks! Audio abstracts available for this article at www.neurosurgery-online.com.
